



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Metolachlor 960

HERBICIDE

ACTIVE CONSTITUENT: 960 g/L METOLACHLOR

GROUP K HERBICIDE

Controls certain annual grasses and broadleaf weeds in certain crops as specified in the directions for use.

CONTENTS
1000 Litres

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

Australis Crop Protection Pty Ltd

ABN 94 150 711 185

Shop 4, 30 Heber Street, Moree NSW 2400

Telephone: 0417 329 133 • Facsimile: 07 3337 9882

www.austcrop.com.au

BATCH NO:

DATE OF MANUFACTURE:



Metolachlor 960

HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Metolachlor 960 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

APVMA Approval No.: 67734/56755

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE



AUSTRALIS
CROP PROTECTION

Australis Crop Protection Pty Ltd

ABN 94 150 711 185

Shop 4, 30 Heber Street, Moree NSW 2400

Telephone: 0417 329 133 • Facsimile: 07 3337 9882

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Metolachlor 960

HERBICIDE

ACTIVE CONSTITUENT: 960 g/L METOLACHLOR

GROUP **K** HERBICIDE

Controls certain annual grasses and broadleaf weeds in certain crops as specified in the directions for use.

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 67734/56755



Australis Crop Protection Pty Ltd
ABN 94 150 711 185
Shop 4, 30 Heber Street, Moree NSW 2400
Telephone: 0417 329 133 • Facsimile: 07 3337 9882
www.austcrop.com.au

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Metolachlor 960 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.



1000L

◀ PEEL HERE ▶

DIRECTIONS FOR USE

CROPS	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Barley, Oats	Suppression of Annual Ryegrass	NSW, ACT, Vic, Tas, SA, WA only	560-750 mL per ha	Apply to moist soils immediately after planting and before crop and weeds emerge. Use the lower rate on light soils. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3 to 4cm of soil should occur within 10 days after spraying. Damage may occur if crop is sown too shallow (less than 4cm). Damage can occur on light sandy soil types after heavy rain. See General Instructions for use in zero-tillage systems.
Wheat, Barley, Oats, Triticale	Toad Rush	NSW, ACT, Vic, Tas, SA, WA only	225 to 375 mL per ha	Apply at or immediately after planting and before crop and weeds emerge. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3 to 4cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required or in situations where crop competition is minimal. When mixed with Triasulfuron 750 apply pre-sowing and incorporate by sowing operation (for wheat only).
Broccoli, Brussels sprouts, Cabbages, Cauliflowers	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	Qld, NSW, ACT, Vic, Tas, NT only	2.25 to 3 litres per ha	Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Blackberry Nightshade, Chickweed, Fat-hen, Lovegrass, Pigeon Grass, Pigweed, Potato Weed and Wireweed are the major problem. See warning – Brassica crops.
	Crowsfoot Grass, Fierce Thornapple (suppression), Liverseed Grass, Winter Grass	NSW, ACT only		Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Apple of Peru, Common Sowthistle, Fierce Thornapple, Liverseed Grass, Redroot Amaranth, Twiggy Turnip and Nettleleaf Goosefoot are the major problems.
	Apple of Peru, Awnless Barnyard Grass, Fierce Thornapple (suppression) Liverseed Grass, Yellow Weed, Redroot Amaranth (suppression)	Qld only		Warning: Qld only, stunting and reduced yield may occur in broccoli in red soils and light alluvial soils.
	Annual Ryegrass, Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Fiddle Dock, Italian Ryegrass, Liverseed Grass, Nettleleaf Goosefoot (suppression), Twiggy Turnip (suppression), Winter Grass	Vic, Tas only		See warning – Brassica crops.

CROPS	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Canola	Toad Rush	Qld, NSW, ACT, Vic, Tas, SA only	225 to 375 mL per ha	Apply at or immediately after planting and before crops and weeds emerge. Apply to moist soil, use rates towards the higher end of the range where longer residual control is required.
	Toad Rush	WA only		Apply before crops and weeds emerge. Apply to moist soil. To avoid crop injury use rates towards the lower end of the range on light sandy soils.
Cotton	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, WA only	1.5 litres per ha	Pre-emergent. Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50% to 70%. See Warning – Cotton.
Green Beans, Navy Beans	Blackberry Nightshade, Common Sowthistle, Fat-hen, Potato Weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass and suppression of Pigweed	Qld, NSW, ACT, Vic, SA only	2.25 to 3 litres per ha	Pre-emergent. Apply before, at or immediately after planting and before crops and weeds have germinated. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control.
	Powell's Amaranth	Tas only	3 litres per ha	Pre-emergent. Apply before, at or immediately after planting and before crops and weeds have germinated. For sufficient Amaranth control the ACP Metolachlor 960 Herbicide application should be followed by 1 L/ha Basagran* (2 leaf Amaranth) plus 2 L/ha Basagran* (4 leaf Amaranth). Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control.

CROPS	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Maize (not Waxy Maize), Sweet Corn	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, ACT, Vic, Tas, SA, NT only	1.5 to 3 litres per hectare	Pre-emergent. Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50% to 70%.
	Blackberry nightshade, Caltrop, Common Thornapple, Mintweed, Needle Burr, Noogoora Burr, Pigweed, Redroot Amaranth, Slim Amaranth		1.5 to 3 litres plus 2.5 litres of Flowable Atrazine (500g/L) per ha	
Pastures – Trikkala Sub-clover, Halfa white clover, Maral persian clover, Paradana Balansa clover, Redquin red clover, Phalaris and Cocksfoot	Toad Rush	NSW, ACT, Vic, SA, NT only	300 to 370 mL per ha	Apply at or immediately after planting and before pasture species and weeds emerge. Application should be made to a moist seed bed. Sufficient rain to thoroughly wet the top 3 to 4cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required. DO NOT use in soils where prolonged waterlogging is anticipated otherwise unacceptable crop retardation will occur. Avoid damage to pasture species by avoiding overlaps and double spraying in headlands where White, Persian and Balansa clovers are sown and if the 500 mL rate is being used.
Peanuts	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew.	Qld, NSW, NT only	1.5 to 3 litres per hectare	Apply before pre-irrigation where soil crusting is likely. Apply before, at, or immediately after planting and before weeds germinate. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm.
Soybeans, Sunflowers		Qld, NSW, ACT, Vic, SA, NT only		DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected.

CROPS	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Sorghum treated with Concep II Sorghum Seed Safener	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, ACT, NT only	1.5 to 3 litres per hectare	Pre-emergent. Apply before, at or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. DO NOT apply to sorghum seed not previously treated with Concep II Sorghum Seed Safener. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4 cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use the lower rate on sandy soils.
	Blackberry Nightshade, Caltrop, Common Thornapple, Mintweed, Needle Burr, Noogoora Burr, Pigweed, Redroot Amaranth, Slim Amaranth		1.5 to 3 litres per hectare plus 2.5 litres of Atrazine Flowable (500g/L) per ha	
Sugar cane (plant and ratoon)	Awnless Barnyard Grass, Barnyard Grass, Bellvine, Billygoat weed (Blue top), Blackberry Nightshade, Caltrop, Common Sowthistle, (Milk Thistle), Common Thornapple, Crowsfoot Grass, Green Summer Grass, Guinea Grass, Mexican Clover, Mintweed, Needle Burr, Passionfruit Vines, Pigeon Grass, Pigweed, Redroot Amaranth, Square Weed, Star of Bethlehem, Summer Grass, Wild Rose	Southern Qld only (Bundaberg South)	1.6 to 2.1 litres plus 2.7 to 3.6 L Atrazine Flowable (500g/L) or 1.5 to 2 kg Atrazine Granules (900g/kg)	Apply as a pre-emergent or post-emergent application to crop. Apply once per year only. Application should be applied to a moist soil before weeds and grasses have germinated. If conditions remain dry for a period of 10 days after spraying, irrigation or shallow cultivation (2.5cm) may assist results. DO NOT throw untreated soil on to treated areas as this will reduce weed control. In Southern Qld use rates toward the higher end of the range on heavy soils or where a high grass population is expected. In Northern Qld, application must be made to moist soil and rainfall or irrigation should occur within 24 hours of application. Use rates towards the higher end of the range where high Green Summer Grass population is expected. Where broadleaf weeds and grasses have emerged and are in the 2 to 4 true leaf stage, knockdown herbicides such as Primatol Z (2 L/ha) and a flowable diuron (500g/L) at 1 L/ha or Paraquat 250 (1.2 to 1.6 L/ha), applied as a post directed spray, can be added. Where broadleaf weeds only have emerged and are at the 2 to 4 true leaf stage then a 500g/L 2,4-D Amine at 2 L/ha should be added. Follow all instructions and restrictions on the Atrazine labels.
		Northern Qld (Mackay North)	2.1 to 2.7 litres plus 3.6 to 4.5 L Atrazine Flowable (500g/L) or 2 to 2.5 kg Atrazine Granules (900g/kg)	
Sweet Potatoes	Black Pigweed, Redshank, Potato Weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrasses, Pigeon Grass, Summer Grass, suppression of Evening Primrose	Qld, NSW, ACT, Vic, SA, NT only	2.25 litres per ha	Pre-emergent. Apply immediately after transplanting before weeds have germinated. Sufficient irrigation to wet the soil through the weed zone should be applied within 24 hours. Further weed germination may occur following rehillng due to exposure of untreated soil.

CROPS	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Tobacco (transplanted in field)	Barnyard Grass, Lovegrass, Pigeon Grass, Summer Grass, and suppression of Common Thornapple (<i>Datura stramonium</i>) and Nightshades (<i>Solanum sarrachoides</i> and <i>S. nigrum</i>)	NSW, Vic only	3 litres per ha	<p>Apply by low volume boom spray prior to transplanting and before weeds and grasses germinate. The seed bed should be fine and free of large clods.</p> <p>Thoroughly incorporate within 7 days of spraying using a rotary hoe or similar equipment set to work to a depth of 7.5cm. DO NOT use disc cultivating implements.</p> <p>An even coverage of the herbicide is essential.</p> <p>The herbicide needs to be activated by moisture through to the top 3 to 4cm of soil at the time of application or within 4 days of incorporation.</p> <p>This can be achieved by applying to moist soil, irrigating soon after incorporation or irrigating immediately after transplanting. Avoid boom overlaps as double treatment may cause growth suppression. Temporary crop retardation may be observed when soil conditions remain dry after application or where black root rot may occur.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

CANOLA: DO NOT GRAZE TREATED FIELDS FOR 10 WEEKS AFTER APPLICATION.

GREEN BEANS: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 10 WEEKS AFTER APPLICATION.

PASTURES: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

SWEET POTATOES: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 23 WEEKS AFTER APPLICATION.

FORAGE ALL OTHER CROPS (EXCEPT FOR SWEET POTATOES): DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

WHEAT, BARLEY, OATS, TRITICALE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

ACP Metolachlor 960 Herbicide is a member of the amide group of herbicides. The product has the diverse sites of mode of action. For weed resistance management this is a Group K herbicide. Some naturally-occurring weed biotypes resistant to ACP Metolachlor 960 Herbicide and other group K herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by ACP Metolachlor 960 Herbicide or other group K herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Australis Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of ACP Metolachlor 960 Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or an Australis Crop Protection representative.

USES

The product is a short residual pre-emergent herbicide for the control of many important annual grasses and broadleaf weeds in the crops nominated.

The product does not control the important broadleaf weeds in maize, sweet corn and Concep II treated sorghum, but when mixed with Flowable Atrazine control of many important annual weeds and grasses is obtained without the problem of carryover of herbicides affecting succeeding crops.

The product acts mainly through shoot (coleoptile) absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the top 3 to 4cm of soil, where it must be present at the time of weed germination.

MIXING

Add the required amount directly to the spray tank, fill with water and mix well.

COMPATIBILITY

ACP Metolachlor 960 Herbicide is compatible with Prometryn 500g/L Flowable Liquid Herbicide, Atrazine 500g/L Flowable Herbicide, Triasulfuron 750WG Herbicide, Paraquat 250g/L and Glyphosate Herbicide.

GROUP K HERBICIDE

WARNING – Cotton: DO NOT use when planting into dry soil or soil with marginal moisture. Under dry planting conditions heavy rain or flood irrigation soon after planting, together with below average soil temperatures, can result in crop injury due to the inability of the germinating seedling to metabolise above normal amounts of herbicide that may accumulate in the root zone as a result of accelerated capillary movement. ACP Metolachlor 960 Herbicide should not be used in situations where these conditions are likely to occur.

WARNING – Brassica crops: Crop retardation may occur where ACP Metolachlor 960 Herbicide is used on soils that have a low organic matter content and which contain more than 60% of fine sand and silt and less than 5% clay in the top 30 cm. On these soil types, use rates of ACP Metolachlor 960 Herbicide towards the lower end of the rate range and apply onto moist soil. Irrigation after application should be limited to no more than 25mm.

Zero – Tillage (Canola, Barley, Oats, Wheat, Triticale): When sowing using knife points and zero-tillage and leaving a seeding slot, DO NOT apply post-seeding unless harrows (or similar) have filled in the seeding slot before application. Alternatively, apply ACP Metolachlor 960 Herbicide immediately before sowing.

DO NOT USE ON SOILS CONTAINING LESS THAN 5% CLAY IN THE TOP 30 cm.

DO NOT USE MORE THAN ONCE PER YEAR.

PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT plant crops other than maize, sweet corn, tobacco, cotton, soybeans, sunflowers, peanuts, broccoli, Brussels sprouts, cabbages, cauliflowers or Concep II treated sorghum within six months of application of ACP Metolachlor 960 Herbicide.

DRIFT WARNING

DO NOT apply under weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

DO NOT graze or cut sweet potato leaves for stock food.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Re-entry Period: DO NOT enter treated area until spray has dried.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Metolachlor 960 Fungicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

* Other trademarks