

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



Quizalofop 200 EC

HERBICIDE

ACTIVE CONSTITUENT: 200 g/L QUIZALOFOP-P-ETHYL
SOLVENTS: 592 g/L HYDROCARBON LIQUID
150 g/L N-METHYL-2-PYRROLIDONE

GROUP A HERBICIDE

For the control of certain grass weeds in broadleaf crops
as per Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 10 LITRES

Australis Crop Protection Pty Ltd
ABN 94 150 711 185
Shop 4, 30 Heber Street, Moree NSW 2400
Ph 0417 329 133 Fax (07) 3337 9882

www.austcrop.com.au

APVMA Approval No: 90180/127911

BATCH NO:

DATE OF MANUFACTURE:

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Quizalofop 200 EC

HERBICIDE

ACTIVE CONSTITUENT: 200 g/L QUIZALOFOP-P-ETHYL
SOLVENTS: 592 g/L HYDROCARBON LIQUID
150 g/L N-METHYL-2-PYRROLIDONE

GROUP **A** HERBICIDE

For the control of certain grass weeds in
broadleaf crops as per Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE OPENING
OR USING THIS PRODUCT

10 LITRES

**UN 3082 ENVIRONMENTALLY
HAZARDOUS SUBSTANCE**

**LIQUID, N.O.S
(CONTAINS
QUIZALOFOP-
P-ETHYL
HAZCHEM
2X
PGIII**

**MISCELLANEOUS
DANGEROUS
GOODS**

9

STORAGE AND DISPOSAL: DO NOT store tank-mixes of ACP Quizalofop 200 EC Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. When storing, keep from contact with fertilisers, fungicides and seeds. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

If the container is a returnable/refillable type, when empty, or contents no longer required, return the container to the point of purchase. This type of container remains the property of Australis Crop Protection Pty Ltd.

SAFETY DIRECTIONS: Harmful if inhaled or swallowed. May irritate the nose and throat. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, mixing, loading, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, and elbow-length chemical resistant gloves. If applying by spraying equipment carried on the back of the user, wear cotton overalls over normal clothing, buttoned to the neck and wrist, and elbow-length chemical resistant gloves. If product on skin immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed do NOT induce vomiting. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Combustible liquid. May be fatal if swallowed and enters airways. Harmful if inhaled. May cause respiratory irritation.

May damage fertility or the unborn child. Very toxic to aquatic life with long lasting effects.

Avoid breathing fumes, mists, vapours or spray. Use only outdoors or in a well ventilated area. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. If skin irritation occurs or eye irritation persists: Get medical advice. Collect spillage.

MATERIAL SAFETY DATA SHEET: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Quizalofop 200 EC Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

APVMA Approval No. 90180/127911



**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

Australis Crop Protection Pty Ltd
ABN 94 150 711 185
Shop 4, 30 Heber Street, Moree NSW 2400
Ph 0417 329 133 Fax (07) 3337 9882

www.austcrop.com.au



▶ PEEL HERE ▶

DIRECTIONS FOR USE**RESTRAINTS:**

DO NOT apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).

DO NOT apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing. Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.

DO NOT apply under conditions of high temperature and low humidity.

DO NOT apply if rainfall is expected within 3 hours.

DO NOT apply less than 50L prepared spray/ha – see section “Ground Spraying”.

BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES.

Crops	Weeds Controlled	Weeds Stage	Rate/ ha	Withholding Period		Critical Comments
				Harvest	Grazing	
Beetroot, Cabbage, Canola, Carrots, Cauliflower, Chickpeas, Clover & subclover pastures, Cucumbers*, Faba beans, Field peas, Honey dew melons*, Lentils, Lucerne, Lupins, Medic, pasture, Mung beans, Navy beans, Onions, Potatoes, Pumpkins*, Radish, Tomatoes, Vetch	Annual Ryegrass (<i>Lolium rigidum</i>) Brome Grasses (<i>Bromus</i> spp.)	3 leaf to early tillering	150mL or 190mL	14 days 9 weeks 11 weeks 10 weeks 14 days 12 weeks	- - 4 weeks - - 4 weeks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Barley grass (<i>Hordeum</i> spp), Volunteer Barley, Volunteer Wheat	3 leaf to early tillering	125mL	- 14 days 12 weeks 9 weeks	- - 4 weeks 4 weeks	Apply when weeds are actively growing*
	Wild Oats (<i>Avena</i> spp)	3 leaf to mid tillering	65mL or 125mL	9 weeks 12 weeks - 6 weeks - 12 weeks 12 weeks 18 weeks 10 weeks 9 weeks 21 days 4 weeks 12 weeks	- 4 weeks 4 weeks 4 weeks 14 days 4 weeks 4 weeks - - - - - - 4 weeks	Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control*

* Always add a surfactant/wetting agent at the recommended rate. See section on “use of surfactant/wetting agent” for details.

- **IMPORTANT: DO NOT** apply ACP Quizalofop 200 EC Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES.

Crops	Weeds Controlled	Weeds Stage	Rate/ ha	Withholding Period		Critical Comments
				Harvest	Grazing	
Beetroot, Cabbage, Cauliflower, Carrots, Cucumbers*, Green beans, Honey dew melons*, Lucerne, Mung beans, Navy beans, Onions, Peanuts, Potatoes, Pineapples, Pumpkins*, Radish, Soybeans, Sunflower, Tomatoes	Awnless Barnyard Grass (<i>Echinochloa colona</i>), Barnyard Grass (<i>Echinochloa crusgalli</i>)	3 leaf to early tillering	250mL or 375mL	14 days 9 weeks 14 weeks 10 weeks 14 days 5 weeks	- - - - - 4 weeks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Couch grass (<i>Cynodon dactylon</i>)	2 – 4 leaf seedling	250mL	9 weeks - 12 weeks 12 weeks 18 weeks 11 weeks 10 weeks 7 days 9 weeks 21 days 12 weeks 9 weeks 4 weeks	- 4 weeks 4 weeks 4 weeks - - - - - - 4 weeks 11 weeks -	Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed*.
	Dinebra (<i>Dinebra retroflexa</i>)	3 leaf to early tillering	125mL or 190mL	18 weeks 11 weeks 10 weeks 9 weeks 21 days 12 weeks 9 weeks 4 weeks	- 11 weeks - - - - - - - 4 weeks 11 weeks -	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Crowsfoot grass (<i>Eleusine indica</i>)	2 leaf to early tillering	250mL	12 weeks 9 weeks 4 weeks	4 weeks 11 weeks -	Apply when weeds are actively growing*.
	Foxtail Millet (<i>Setaria italica</i>)	2 leaf to early tillering	125mL or 250mL			Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Columbus grass (<i>Sorghum x almum</i>), Johnson grass (<i>Sorghum halepense</i>), Liverseed grass (<i>Urochloa panicoides</i>)	2 leaf to 3 tillering	250mL or 375mL			Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Rhodes grass (<i>Chloris gayana</i>), Stink grass (<i>Eragrostis ciliaris</i>)	3 leaf to early tillering	125mL or 250mL			Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Queensland Blue grass (<i>Dichanthium sericeum</i>)	3 leaf to early tillering	250mL			Apply when weeds are actively growing*.
	Summer grass (<i>Digitaria ciliaris</i>)	3 leaf to early tillering	250mL or 500mL			Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*

* Always add a surfactant/wetting agent at the recommended rate. See section on “use of surfactant/wetting agent” for details.

- **IMPORTANT: DO NOT** apply ACP Quizalofop 200 EC Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

HAND GUN SPRAYING: ALL STATES

Crops	Weeds Controlled	Weeds Stage	Rate/ha	Withholding Period		Critical Comments
				Harvest	Grazing	
Beetroot, Cabbage, Cauliflower, Canola, Carrots, Chickpeas, Clover & subclover pastures, Cucumbers•, Faba beans, Field peas, Grapes, Green beans, Honey dew melons•, Lentils, Lucerne, Lupins, Medic pasture, Mung beans, Navy beans, Non crop, Onions, Peanuts, Pineapples, Potatoes, Pumpkins•, Radish, Soy beans, Sunflowers, Tomatoes, Vetch	Awnless BarnyardGrass (<i>Echinochloa colona</i>), Crowsfoot grass (<i>Eleusine indica</i>)	3 leaf to early tillering	65 mL	14 days	-	Apply when weeds are actively growing. Use a minimum of 800L of prepared spray/ha. Thoroughly wet target weeds (especially well-established clumps) without causing runoff. Repeat spraying may be necessary for perennial grass species. Always add a surfactant/wetting agent at the recommended rate – see section on “Use of Surfactant/Wetting agent” for details.
				9 weeks	-	
				14 days	-	
				11 weeks	4 weeks	
				10 weeks	-	
				12 weeks	4 weeks	
	-	14 days				
	Paspalum (<i>Paspalum dilatatum</i>)	Seedling to Established Clumps	14 days	-		
				12 weeks	4 weeks	
	Johnson grass (<i>Sorghum halepense</i>)	Seedling 2 leaf to early tillering	9 weeks	4 weeks		
				-		
	Rhizome	2 leaf to early tillering	5 weeks	4 weeks		
9 weeks				-		
Kikuyu (<i>Pennisetum clandestinum</i>)	Seedling to Established plants	125 mL	12 weeks	4 weeks		
			-	4 weeks		
			6 weeks	4 weeks		
			-	14 days		
			12 weeks	4 weeks		
			12 weeks	4 weeks		
			-	-		
			18 weeks	-		
			11 weeks	11 weeks		
			7 days	-		
10 weeks	-					
9 weeks	-					
21 days	-					
12 weeks	4 weeks					
9 weeks	11 weeks					
4 weeks	-					
12 weeks	4 weeks					

- **IMPORTANT: DO NOT** apply ACP Quizalofop 200 EC Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

HARVEST

BEETROOT, CAULIFLOWER, CUCUMBERS:

DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

CABBAGE, FIELD PEAS, HONEY DEW MELON, PUMPKINS, SUNFLOWERS:

DO NOT HARVEST FOR 9 WEEKS AFTER APPLICATION.

CARROTS, POTATOES:

DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

CHICKPEAS, FABA BEANS, LENTILS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH:

DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION.

CANOLA, PEANUTS:

DO NOT HARVEST FOR 11 WEEKS AFTER APPLICATION.

GREEN BEANS:

DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

GRAPES:

NOT REQUIRED WHEN USED AS DIRECTED.

LUPINS:

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

ONIONS:

DO NOT HARVEST FOR 18 WEEKS AFTER APPLICATION.

PINEAPPLES:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

RADISH:

DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

TOMATOES:

DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

GRAZING

CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, GREEN BEANS, LENTILS, LUCERNE, LUPINS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH:

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION.

CLOVER, MEDIC PASTURES (EXCEPT LUCERNE):

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.

PEANUTS, SUNFLOWERS: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 11 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

ACP Quizalofop 200 EC Herbicide is an emulsifiable concentrate herbicide used post emergence for the control of certain grass weeds in specified broadleaf crops.

ACP Quizalofop 200 EC Herbicide is a systemic herbicide, which is rapidly absorbed and readily translocates from the treated foliage to the root system and growing points of the plant. Applications made to grass weeds stressed due to insufficient moisture or cold temperatures may result in unsatisfactory control.

Time of Application

For maximum benefit, weeds should be sprayed when young and actively growing. Application may be made at any stage of crop growth (except to cucurbit crops), but sufficient time must be allowed for the prescribed withholding period to expire before harvest and / or grazing or cutting for stockfeed.

IMPORTANT: DO NOT apply ACP Quizalofop 200 EC Herbicide to cucurbit crops before the five true leaf stage.

Spray Preparation

If spray equipment has been previously used for the application of sulfonylurea herbicides e.g. (Metsulfuron methyl), chlorsulfuron, thifensulfuron methyl, triasulfuron, ensure that the recommended sprayer equipment procedure for the product previously applied has been followed, before using the equipment for the application of ACP Quizalofop 200 EC Herbicide. ACP Quizalofop 200 EC Herbicide is an emulsifiable concentrate to be mixed with water and applied as a spray. Add the required amount of ACP Quizalofop 200 EC Herbicide directly to the spray tank and mix well. If mixing with mancozeb or carbendazim, add these to the spray tank and mix thoroughly PRIOR to adding ACP Quizalofop 200 EC Herbicide. If spray solution has been left standing for more than 3 hours ensure thorough re-agitation before commencement of spraying. DO NOT allow spray mixture to stand overnight, as re-suspension may be difficult.

Use of a Surfactant/Wetting Agent

ACP Quizalofop 200 EC Herbicide may be used with either of the following adjuvant system:

- 1) A methylated and/or ethylated seed crop oil (e.g. Hasten#) at 1L/100L of final spray volume (i.e. 1% v/v).
- 2) A non-ionic surfactant (1000 gai/L) at 200 mL/100L of final spray volume (i.e. 0.2% v/v).

DO NOT use ACP Quizalofop 200 EC Herbicide with mineral based crop oils or non-methylated spray oils of crop origin (e.g. Codacide# oil, Synerrol# oil) as spray adjuvants.

Compatibilities

ACP Quizalofop 200 EC Herbicide is compatible with the herbicides sethoxydim and clethodim. The recommended surfactant/wetting agent is the methylated and/or ethylated seed crop oil adjuvant system. ACP Quizalofop 200 EC Herbicide is compatible with the insecticides deltamethrin, omethoate, dimethoate, methomyl, endosulfan, the fungicides carbendazim and mancozeb and the herbicide clopyralid when using the methylated and/or ethylated seed crop oil adjuvant system. ACP Quizalofop 200 EC Herbicide is compatible with the herbicide diflufenican when using the non-ionic surfactant only adjuvant system. DO NOT add crop oil to tank mixtures of ACP Quizalofop

200 EC Herbicide /diflufenican. Application of ACP Quizalofop 200 EC Herbicide /diflufenican tank mix may result in crop effects, which usually appear as yellow or white bleaching on the leaves. New growth and subsequent yield, however, will not be affected.

To minimise crop effects apply ACP Quizalofop 200 EC Herbicide a minimum of 5 days prior to diflufenican application or a minimum 10 days after diflufenican application. (14-21 days if weeds have been stressed).

Ground Spraying

Use between 50 and 100 litres of prepared spray/ha, the higher spray volume gives more reliable results. If weeds are large and the crop canopy is dense, increase the volume to 150 litres prepared spray/ha.

Use sprayers with flat fan or hollow cone nozzles that produce droplets within the FINE-MEDIUM droplet size category. 1100 flat fan nozzles are preferred. For proper spray coverage adjust the boom and nozzle height according to the specification listed by the nozzle manufacturers. Ensure the spray is evenly distributed and a double overlap spray pattern is obtained. Avoid overlapping boomspray widths and shut off spray booms while starting, slowing or stopping.

Sprayer cleanup

Thoroughly clean all traces of ACP Quizalofop 200 EC Herbicide from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean those parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed cereal and grass crops.

If spray equipment has been previously used for the application of sulfonyleurea herbicides e.g. (Metsulfuron methyl), thifensulfuron methyl, chlorsulfuron, triasulfuron ensure that the recommended sprayer cleanup procedure of the product previously sprayed has been followed before using the equipment for the application of ACP Quizalofop 200 EC Herbicide.

GROUP A HERBICIDE

RESISTANT WEEDS WARNING

ACP Quizalofop 200 EC Herbicide is a member of the "Fop" (aryloxyphenoxy propionate) group of herbicides. ACP Quizalofop 200 EC Herbicide has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management ACP Quizalofop 200 EC Herbicide is a Group A herbicide. Some naturally occurring weed biotypes resistant to ACP Quizalofop 200 EC Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by ACP Quizalofop 200 EC Herbicide or other Group A herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Australis Crop Protection Pty Ltd accepts no liability for any losses that may result from failure of ACP Quizalofop 200 EC Herbicide to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, Australis Crop Protection Pty Ltd Representative, local Department of Agriculture or Primary Industries.

RE-ENTRY

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of ACP Quizalofop 200 EC Herbicide.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

(10L, 20L and 200L only)

DO NOT store tank-mixes of ACP Quizalofop 200 EC Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. When storing, keep from contact with fertilisers, fungicides and seeds. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

If the container is a returnable/refillable type, when empty, or contents no longer required, return the container to the point of purchase. This type of container remains the property of Australis Crop Protection Pty Ltd.

Refillable Containers:

(1000L only)

Store in closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container:

Store the original sealed ENVIRODRUM in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with Micro Matic valve or security seal. DO NOT contaminate the ENVIRODRUM with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the ENVIRODRUM have been used, please return the empty ENVIRODRUM to the point of purchase. The ENVIRODRUM remains the property of Australis Crop Protection Pty Ltd.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. May irritate the nose and throat. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, mixing, loading, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, and elbow-length chemical resistant gloves. If applying by spraying equipment carried on the back of the user, wear cotton overalls over normal clothing, buttoned to the neck and wrist, and elbow-length chemical resistant gloves. If product on skin immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed do NOT induce vomiting. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

CONDITIONS OF SALE

The use of ACP Quizalofop 200 EC Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Not a trademark of Australis Crop Protection Pty Ltd.





Quizalofop 200 EC

HERBICIDE

STORAGE AND DISPOSAL: DO NOT store tank-mixes of ACP Quizalofop 200 EC Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. When storing, keep from contact with fertilisers, fungicides and seeds. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

If the container is a returnable/refillable type, when empty, or contents no longer required, return the container to the point of purchase. This type of container remains the property of Australis Crop Protection Pty Ltd.

SAFETY DIRECTIONS: Harmful if inhaled or swallowed. May irritate the nose and throat. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, mixing, loading, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, and elbow-length chemical resistant gloves. If applying by spraying equipment carried on the back of the user, wear cotton overalls over normal clothing, buttoned to the neck and wrist, and elbow-length chemical resistant gloves. If product on skin immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed do NOT induce vomiting. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Combustible liquid. May be fatal if swallowed and enters airways. Harmful if inhaled. May cause respiratory irritation. May damage fertility or the unborn child. Very toxic to aquatic life with long lasting effects.

Avoid breathing fumes, mists, vapours or spray. Use only outdoors or in a well ventilated area. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. If skin irritation occurs or eye irritation persists: Get medical advice. Collect spillage.

MATERIAL SAFETY DATA SHEET: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Quizalofop 200 EC Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

APVMA Approval No. 90180/127911

UN 3082
ENVIRONMENTALLY HAZARDOUS
SUBSTANCE LIQUID, N.O.S
(CONTAINS
QUIZALOFOP-
P-ETHYL
HAZCHEM
2X
PGIII

MISCELLANEOUS
DANGEROUS
GOODS

9



IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE